

1. Compute the sum of all the roots of  $(2x + 3)(x - 4) + (2x + 3)(x - 6) = 0$ .

- (A)  $7/2$     (B) 4    (C) 5    (D) 7    (E) 13

2. Cindy was asked by her teacher to subtract 3 from a certain number and then divide the result by 9. Instead, she subtracted 9 and then divided the result by 3, giving an answer of 43. What would her answer have been had she worked the problem correctly?

- (A) 15    (B) 34    (C) 43    (D) 51    (E) 138

3. According to the standard convention for exponentiation,

$$2^{2^{2^2}} = 2^{\left(2^{(2^2)}\right)} = 2^{16} = 65,536.$$


If the order in which the exponentiations are performed is changed, how many other values are possible?

- (A) 0    (B) 1    (C) 2    (D) 3    (E) 4

4. Find the degree measure of an angle whose complement is 25% of its supplement.

- (A) 48    (B) 60    (C) 75    (D) 120    (E) 150

5. Each of the small circles in the figure has radius one. The innermost circle is tangent to the six circles that surround it, and each of those circles is tangent to the large circle and to its small-circle neighbors. Find the area of the shaded region.



- (A)  $\pi$     (B)  $1.5\pi$     (C)  $2\pi$     (D)  $3\pi$     (E)  $3.5\pi$


6. For how many positive integers  $m$  does there exist at least one positive integer  $n$  such that  $m \cdot n \leq m + n$ ?

- (A) 4    (B) 6    (C) 9    (D) 12    (E) infinitely many

7. If an arc of  $45^\circ$  on circle  $A$  has the same length as an arc of  $30^\circ$  on circle  $B$ , then the ratio of the area of circle  $A$  to the area of circle  $B$  is

(A)  $\frac{4}{9}$     (B)  $\frac{2}{3}$     (C)  $\frac{5}{6}$     (D)  $\frac{3}{2}$     (E)  $\frac{9}{4}$

8. Betsy designed a flag using blue triangles ( $\blacktriangle$ ), small white squares ( $\square$ ), and a red center square ( $\blacksquare$ ), as shown. Let  $B$  be the total area of the blue triangles,  $W$  the total area of the white squares, and  $R$  the area of the red square. Which of the following is correct?



(A)  $B = W$     (B)  $W = R$     (C)  $B = R$     (D)  $3B = 2R$     (E)  $2R = W$

9. Jamal wants to store 30 computer files on floppy disks, each of which has a capacity of 1.44 megabytes (mb). Three of his files require 0.8 mb of memory each, 12 more require 0.7 mb each, and the remaining 15 require 0.4 mb each. No file can be split between floppy disks. What is the minimal number of floppy disks that will hold all the files?
- (A) 12    (B) 13    (C) 14    (D) 15    (E) 16
10. Sarah pours four ounces of coffee into an eight-ounce cup and four ounces of cream into a second cup of the same size. She then transfers half the coffee from the first cup to the second and, after stirring thoroughly, transfers half the liquid in the second cup back to the first. What fraction of the liquid in the first cup is now cream?
- (A)  $\frac{1}{4}$     (B)  $\frac{1}{3}$     (C)  $\frac{3}{8}$     (D)  $\frac{2}{5}$     (E)  $\frac{1}{2}$
11. Mr. Earl E. Bird leaves his house for work at exactly 8:00 A.M. every morning. When he averages 40 miles per hour, he arrives at his workplace three minutes late. When he averages 60 miles per hour, he arrives three minutes early. At what average speed, in miles per hour, should Mr. Bird drive to arrive at his workplace precisely on time?
- (A) 45    (B) 48    (C) 50    (D) 55    (E) 58
12. Both roots of the quadratic equation  $x^2 - 63x + k = 0$  are prime numbers. The number of possible values of  $k$  is
- (A) 0    (B) 1    (C) 2    (D) 4    (E) more than four

13. Two different positive numbers  $a$  and  $b$  each differ from their reciprocals by 1. What is  $a + b$ ?

(A) 1    (B) 2    (C)  $\sqrt{5}$     (D)  $\sqrt{6}$     (E) 3

14. For all positive integers  $n$ , let  $f(n) = \log_{2002} n^2$ . Let

$$N = f(11) + f(13) + f(14).$$

Which of the following relations is true?

(A)  $N > 1$     (B)  $N = 1$     (C)  $1 < N < 2$     (D)  $N = 2$     (E)  $N > 2$

15. The mean, median, unique mode, and range of a collection of eight integers are all equal to 8. The largest integer that can be an element of this collection is

(A) 11    (B) 12    (C) 13    (D) 14    (E) 15

16. Tina randomly selects two distinct numbers from the set  $\{1, 2, 3, 4, 5\}$ , and Sergio randomly selects a number from the set  $\{1, 2, \dots, 10\}$ . The probability that Sergio's number is larger than the sum of the two numbers chosen by Tina is

(A)  $2/5$     (B)  $9/20$     (C)  $1/2$     (D)  $11/20$     (E)  $24/25$

17. Several sets of prime numbers, such as  $\{7, 83, 421, 659\}$ , use each of the nine nonzero digits exactly once. What is the smallest possible sum such a set of primes could have?

(A) 193    (B) 207    (C) 225    (D) 252    (E) 477

18. Let  $C_1$  and  $C_2$  be circles defined by

$$(x - 10)^2 + y^2 = 36$$


and

$$(x + 15)^2 + y^2 = 81,$$


respectively. What is the length of the shortest line segment  $\overline{PQ}$  that is tangent to  $C_1$  at  $P$  and to  $C_2$  at  $Q$ ?

(A) 15    (B) 18    (C) 20    (D) 21    (E) 24

19. The graph of the function  $f$  is shown below. How many solutions does the equation  $f(f(x)) = 6$  have?



- (A) 2    (B) 4    (C) 5    (D) 6    (E) 7
20. Suppose that  $a$  and  $b$  are digits, not both nine and not both zero, and the repeating decimal  $0.\overline{ab}$  is expressed as a fraction in lowest terms. How many different denominators are possible?
- (A) 3    (B) 4    (C) 5    (D) 8    (E) 9
21. Consider the sequence of numbers: 4, 7, 1, 8, 9, 7, 6,  $\dots$ . For  $n > 2$ , the  $n$ th term of the sequence is the units digit of the sum of the two previous terms. Let  $S_n$  denote the sum of the first  $n$  terms of this sequence. The smallest value of  $n$  for which  $S_n > 10,000$  is:
- (A) 1992    (B) 1999    (C) 2001    (D) 2002    (E) 2004
22. Triangle  $ABC$  is a right triangle with  $\angle ACB$  as its right angle,  $m\angle ABC = 60^\circ$ , and  $AB = 10$ . Let  $P$  be randomly chosen inside  $\triangle ABC$ , and extend  $\overline{BP}$  to meet  $\overline{AC}$  at  $D$ . What is the probability that  $BD > 5\sqrt{2}$ ?



- (A)  $\frac{2 - \sqrt{2}}{2}$     (B)  $\frac{1}{3}$     (C)  $\frac{3 - \sqrt{3}}{3}$     (D)  $\frac{1}{2}$     (E)  $\frac{5 - \sqrt{5}}{5}$


23. In triangle  $ABC$ , side  $\overline{AC}$  and the perpendicular bisector of  $\overline{BC}$  meet in point  $D$ , and  $\overline{BD}$  bisects  $\angle ABC$ . If  $AD = 9$  and  $DC = 7$ , what is the area of triangle  $ABD$ ?

- (A) 14    (B) 21    (C) 28    (D)  $14\sqrt{5}$     (E)  $28\sqrt{5}$

24. Find the number of ordered pairs of real numbers  $(a, b)$  such that  $(a+bi)^{2002} = a - bi$ .

- (A) 1001    (B) 1002    (C) 2001    (D) 2002    (E) 2004

25. The nonzero coefficients of a polynomial  $P$  with real coefficients are all replaced by their mean to form a polynomial  $Q$ . Which of the following could be a graph of  $y = P(x)$  and  $y = Q(x)$  over the interval  $-4 \leq x \leq 4$ ?



Solutions

1. (A) Factor to get  $(2x + 3)(2x - 10) = 0$ , so the two roots are  $-3/2$  and  $5$ , which sum to  $7/2$ .

2. (A) Let  $x$  be the number she was given. Her calculations produce

$$\frac{x - 9}{3} = 43,$$

so

$$x - 9 = 129 \quad \text{and} \quad x = 138.$$

The correct answer is

$$\frac{138 - 3}{9} = \frac{135}{9} = 15.$$

3. (B) No matter how the exponentiations are performed,  $2^{2^2}$  always gives 16. Depending on which exponentiation is done last, we have

$$(2^{2^2})^2 = 256, \quad 2^{(2^{2^2})} = 65,536, \quad \text{or} \quad (2^2)^{(2^2)} = 256,$$

so there is one other possible value.

4. (B) The appropriate angle  $x$  satisfies

$$90 - x = \frac{1}{4}(180 - x), \quad \text{so} \quad 360 - 4x = 180 - x.$$

Solving for  $x$  gives  $3x = 180$ , so  $x = 60$ .

5. (C) The large circle has radius 3, so its area is  $\pi \cdot 3^2 = 9\pi$ . The seven small circles have a total area of  $7(\pi \cdot 1^2) = 7\pi$ . So the shaded region has area  $9\pi - 7\pi = 2\pi$ .

6. (E) When  $n = 1$ , the inequality becomes  $m \leq 1 + m$ , which is satisfied by all integers  $m$ . Thus, there are infinitely many of the desired values of  $m$ .

7. (A) Let  $C_A = 2\pi R_A$  be the circumference of circle  $A$ , let  $C_B = 2\pi R_B$  be the circumference of circle  $B$ , and let  $L$  the common length of the two arcs. Then

$$\frac{45}{360}C_A = L = \frac{30}{360}C_B.$$

Solutions


Therefore

$$\frac{C_A}{C_B} = \frac{2}{3} \quad \text{so} \quad \frac{2}{3} = \frac{2\pi R_A}{2\pi R_B} = \frac{R_A}{R_B}.$$

Thus, the ratio of the areas is

$$\frac{\text{Area of Circle (A)}}{\text{Area of Circle (B)}} = \frac{\pi R_A^2}{\pi R_B^2} = \left(\frac{R_A}{R_B}\right)^2 = \frac{4}{9}.$$

8. (A) Draw additional lines to cover the entire figure with congruent triangles. There are 24 triangles in the blue region, 24 in the white region, and 16 in the red region. Thus,  $B = W$ .



9. (B) First note that the amount of memory needed to store the 30 files is

$$3(0.8) + 12(0.7) + 15(0.4) = 16.8 \text{ mb},$$

so the number of disks is at least

$$\frac{16.8}{1.44} = 11 + \frac{2}{3}.$$

However, a disk that contains a 0.8-mb file can, in addition, hold only one 0.4-mb file, so on each of these disks at least 0.24 mb must remain unused. Hence, there is at least  $3(0.24) = 0.72$  mb of unused memory, which is equivalent to half a disk. Since

$$\left(11 + \frac{2}{3}\right) + \frac{1}{2} > 12,$$

at least 13 disks are needed.

To see that 13 disks suffice, note that:

Six disks could be used to store the 12 files containing 0.7 mb;

Three disks could be used to store the three 0.8-mb files together with three of the 0.4-mb files;

Four disks could be used to store the remaining twelve 0.4-mb files.

### Solutions

10. (D) After the first transfer, the first cup contains two ounces of coffee, and the second cup contains two ounces of coffee and four ounces of cream. After the second transfer, the first cup contains  $2 + (1/2)(2) = 3$  ounces of coffee and  $(1/2)(4) = 2$  ounces of cream. Therefore, the fraction of the liquid in the first cup that is cream is  $2/(2 + 3) = 2/5$ .

11. (B) Let  $t$  be the number of hours Mr. Bird must travel to arrive on time. Since three minutes is the same as 0.05 hours,  $40(t + 0.05) = 60(t - 0.05)$ . Thus,

$$40t + 2 = 60t - 3, \quad \text{so } t = 0.25.$$

The distance from his home to work is  $40(0.25 + 0.05) = 12$  miles. Therefore, his average speed should be  $12/0.25 = 48$  miles per hour.

OR

Let  $d$  be the distance from Mr. Bird's house to work, and let  $s$  be the desired average speed. Then the desired driving time is  $d/s$ . Since  $d/60$  is three minutes too short and  $d/40$  is three minutes too long, the desired time must be the average, so

$$\frac{d}{s} = \frac{1}{2} \left( \frac{d}{60} + \frac{d}{40} \right).$$

This implies that  $s = 48$ .

12. (B) Let  $p$  and  $q$  be two primes that are roots of  $x^2 - 63x + k = 0$ . Then

$$x^2 - 63x + k = (x - p)(x - q) = x^2 - (p + q)x + p \cdot q,$$

so  $p + q = 63$  and  $p \cdot q = k$ . Since 63 is odd, one of the primes must be 2 and the other 61. Thus, there is exactly one possible value for  $k$ , namely  $k = p \cdot q = 2 \cdot 61 = 122$ .

13. (C) A number  $x$  differs by one from its reciprocal if and only if  $x - 1 = 1/x$  or  $x + 1 = 1/x$ . These equations are equivalent to  $x^2 - x - 1 = 0$  and  $x^2 + x - 1 = 0$ . Solving these by the quadratic formula yields the positive solutions

$$\frac{1 + \sqrt{5}}{2} \quad \text{and} \quad \frac{-1 + \sqrt{5}}{2},$$

which are reciprocals of each other. The sum of the two numbers is  $\sqrt{5}$ .



Solutions

14. (D) We have

$$N = \log_{2002} 11^2 + \log_{2002} 13^2 + \log_{2002} 14^2 = \log_{2002} 11^2 \cdot 13^2 \cdot 14^2 = \log_{2002} (11 \cdot 13 \cdot 14)^2.$$

Simplifying gives

$$N = \log_{2002} (11 \cdot 13 \cdot 14)^2 = \log_{2002} 2002^2 = 2.$$

15. (D) The values 6, 6, 6, 8, 8, 8, 8, 14 satisfy the requirements of the problem, so the answer is at least 14. If the largest number were 15, the collection would have the ordered form 7,  $\underline{\quad}$ ,  $\underline{\quad}$ , 8, 8,  $\underline{\quad}$ ,  $\underline{\quad}$ , 15. But  $7+8+8+15 = 38$ , and a mean of 8 implies that the sum of all values is 64. In this case, the four missing values would sum to  $64 - 38 = 26$ , and their average value would be 6.5. This implies that at least one would be less than 7, which is a contradiction. Therefore, the largest integer that can be in the set is 14.


16. (A) There are ten ways for Tina to select a pair of numbers. The sums 9, 8, 4, and 3 can be obtained in just one way, and the sums 7, 6, and 5 can each be obtained in two ways. The probability for each of Sergio's choices is  $1/10$ . Considering his selections in decreasing order, the total probability of Sergio's choice being greater is

$$\left(\frac{1}{10}\right) \left(1 + \frac{9}{10} + \frac{8}{10} + \frac{6}{10} + \frac{4}{10} + \frac{2}{10} + \frac{1}{10} + 0 + 0 + 0\right) = \frac{2}{5}.$$

17. (B) First, observe that 4, 6, and 8 cannot be the units digit of any two-digit prime, so they must contribute at least  $40+60+80 = 180$  to the sum. The remaining digits must contribute at least  $1+2+3+5+7+9 = 27$  to the sum. Thus, the sum must be at least 207, and we can achieve this minimum only if we can construct a set of three one-digit primes and three two-digit primes. Using the facts that nine is not prime and neither two nor five can be the units digit of any two-digit prime, we can construct the sets  $\{2, 3, 5, 41, 67, 89\}$ ,  $\{2, 3, 5, 47, 61, 89\}$ , or  $\{2, 5, 7, 43, 61, 89\}$ , each of which yields a sum of 207.

Solutions

18. (C) The centers are at  $A = (10, 0)$  and  $B = (-15, 0)$ , and the radii are 6 and 9, respectively. Since the internal tangent is shorter than the external tangent,  $\overline{PQ}$  intersects  $\overline{AB}$  at a point  $D$  that divides  $\overline{AB}$  into parts proportional to the radii. The right triangles  $\triangle APD$  and  $\triangle BQD$  are similar with ratio of similarity 2 : 3. Therefore,  $D = (0, 0)$ ,  $PD = 8$ , and  $QD = 12$ . Thus  $PQ = 20$ .



19. (D) The equation  $f(f(x)) = 6$  implies that  $f(x) = -2$  or  $f(x) = 1$ . The horizontal line  $y = -2$  intersects the graph of  $f$  twice, so  $f(x) = -2$  has two solutions. Similarly,  $f(x) = 1$  has 4 solutions, so there are 6 solutions of  $f(f(x)) = 6$ .
20. (C) Since  $0.\overline{ab} = \frac{ab}{99}$ , the denominator must be a factor of  $99 = 3^2 \cdot 11$ . The factors of 99 are 1, 3, 9, 11, 33, and 99. Since  $a$  and  $b$  are not both nine, the denominator cannot be 1. By choosing  $a$  and  $b$  appropriately, we can make fractions with each of the other denominators.
21. (B) Writing out more terms of the sequence yields

4, 7, 1, 8, 9, 7, 6, 3, 9, 2, 1, 3, 4, 7, 1...


The sequence repeats itself, starting with the 13th term. Since  $S_{12} = 60$ ,  $S_{12k} = 60k$  for all positive integers  $k$ . The largest  $k$  for which  $S_{12k} \leq 10,000$  is

$$k = \left\lfloor \frac{10,000}{60} \right\rfloor = 166,$$

and  $S_{12 \cdot 166} = 60 \cdot 166 = 9960$ . To have  $S_n > 10,000$ , we need to add enough additional terms for their sum to exceed 40. This can be done by adding the next 7 terms of the sequence, since their sum is 42. Thus, the smallest value of  $n$  is  $12 \cdot 166 + 7 = 1999$ .


22. (C) Since  $AB$  is 10, we have  $BC = 5$  and  $AC = 5\sqrt{3}$ . Choose  $E$  on  $\overline{AC}$  so that  $CE = 5$ . Then  $BE = 5\sqrt{2}$ . For  $BD$  to be greater than  $5\sqrt{2}$ ,  $P$  has to be inside  $\triangle ABE$ . The probability that  $P$  is inside  $\triangle ABE$  is

$$\frac{\text{Area of } \triangle ABE}{\text{Area of } \triangle ABC} = \frac{\frac{1}{2}EA \cdot BC}{\frac{1}{2}CA \cdot BC} = \frac{EA}{AC} = \frac{5\sqrt{3} - 5}{5\sqrt{3}} = \frac{\sqrt{3} - 1}{\sqrt{3}} = \frac{3 - \sqrt{3}}{3}.$$



23. (D) By the angle-bisector theorem,  $\frac{AB}{BC} = \frac{9}{7}$ . Let  $AB = 9x$  and  $BC = 7x$ , let  $m\angle ABD = m\angle CBD = \theta$ , and let  $M$  be the midpoint of  $\overline{BC}$ . Since  $M$  is on the perpendicular bisector of  $\overline{BC}$ , we have  $BD = DC = 7$ . Then

$$\cos \theta = \frac{\frac{7x}{2}}{7} = \frac{x}{2}.$$



Applying the Law of Cosines to  $\triangle ABD$  yields

$$9^2 = (9x)^2 + 7^2 - 2(9x)(7) \left(\frac{x}{2}\right),$$

from which  $x = 4/3$  and  $AB = 12$ . Apply Heron's formula to obtain the area of triangle  $ABD$  as  $\sqrt{14 \cdot 2 \cdot 5 \cdot 7} = 14\sqrt{5}$ .

24. (E) Let  $z = a + bi$ ,  $\bar{z} = a - bi$ , and  $|z| = \sqrt{a^2 + b^2}$ . The given relation becomes  $z^{2002} = \bar{z}$ . Note that

$$|z|^{2002} = |z^{2002}| = |\bar{z}| = |z|,$$

24. (E) Let  $z = a + bi$ ,  $\bar{z} = a - bi$ , and  $|z| = \sqrt{a^2 + b^2}$ . The given relation becomes  $z^{2002} = \bar{z}$ . Note that

$$|z|^{2002} = |z^{2002}| = |\bar{z}| = |z|,$$

from which it follows that

$$|z| (|z|^{2001} - 1) = 0.$$

Hence  $|z| = 0$ , and  $(a, b) = (0, 0)$ , or  $|z| = 1$ . In the case  $|z| = 1$ , we have  $z^{2002} = \bar{z}$ , which is equivalent to  $z^{2003} = \bar{z} \cdot z = |z|^2 = 1$ . Since the equation  $z^{2003} = 1$  has 2003 distinct solutions, there are altogether  $1 + 2003 = 2004$  ordered pairs that meet the required conditions.

25. (B) The sum of the coefficients of  $P$  and the sum of the coefficients of  $Q$  will be equal, so  $P(1) = Q(1)$ . The only answer choice with an intersection at  $x = 1$  is (B). (The polynomials in graph B are  $P(x) = 2x^4 - 3x^2 - 3x - 4$  and  $Q(x) = -2x^4 - 2x^2 - 2x - 2$ .)

# 2002 ANSWERS

Question	Correct Answer
1	A
2	A
3	B
4	B
5	C
6	E
7	A
8	A
9	B
10	D
11	B
12	B
13	C
14	D
15	D
16	A
17	B
18	C
19	D
20	C
21	B
22	C
23	D
24	E
25	B